AIRS RT model comparison



R. Saunders with inputs from

V. Sherlock, A. Von Engeln, N. Bormann, S. Hannon, J-L Moncet, Y. Han, D.S. Turner

Activity

- Compare AIRS RT models
- channels for 52 profiles
- For some models compute jacobians for a selection of ~100 channels
- Document run times
- n Complete by Jan 04

Participants and status

lel	Participant	Dire
V-7	R. Saunders, METO	Yes
an [¶]	Y. Han, NESDIS	Yes
S	J-L. Moncet, AER	Yes
TM	J-L. Moncet, AER	Yes
M	N. Bormann, ECMWF	Yes
ppod	V. Sherlock, NIWA	Yes
rs	A. VEngeln, Bremen	Yes
TA	S. Hannon, UMBC	Yes
	W. Smith, NASA	
	S. Heilliette, LMD	
3L	D.S. Turner, MSC	Yes
AST	L. Garand, MSC	
SI	C. Serio, Uni Bas	
ode	F. Miskolczi, NASA	


Mod


LBL


ISC

Initial results

- To date have only compared direct calcs for 9 models who have submitted results
- n Used RFM as reference model (this may favour models based on GENLN2)
- Bias and sdev plots shown of differences for each AIRS channel for 51 diverse profiles










AIRS RT model comparison (2)









Summary

- n Different models have mean differences ~0.1K in some spectral regions
- n Some differences match those seen in LIE comparison
- nIn a few spectral regions differences can be up to 0.3K for some models
- n Against RFM all fast models show a reasonable performance but there are differences
- n No work done on Jacobian results yet
- n More participants encouraged; Plan is to report in Spring 2004

International TOVS Study Conference, 13th, TOVS-13, Sainte Adele, Quebec, Canada, 29 October-4 November 2003. Madison, WI, University of Wisconsin-Madison, Space Science and Engineering Center, Cooperative Institute for Meteorological Satellite Studies, 2003.