# CURRENT AND FUTURE SATELLITE PROGRAMS AND SYSTEMS IN INDIA

Devendra Singh and R.C.Bhatia Email: dschahar@hotmail.com

# INDIA METEOROLOGICAL DEPARTMENT NEW DELHI, INDIA

## **INSAT Meteorological Satellite Applications Programme**

- Round the clock surveillance of weather systems including severe weather events around the Indian region.
- Operational parameters for weather forecasting cloud cover, cloud top temperature, sea surface temperature, snow cover, cloud motion vector, outgoing long wave radiation etc.
- Collection and transmission of meteorological, hydrological and oceanographic data from remote/inaccessible areas through Data Collection Platforms.
- Timely dissemination of warning of impending disasters such as cyclones through Cyclone Warning Dissemination Systems.
- Dissemination of Meteorological information including processed images of weather systems through Meteorological Data dissemination system

## INDIAN NATIONAL SATELLITE (INSAT) - PAST & PRESENT

Satellite	Lunch date	Met Payload with Wavelength Bands	Major Applications	Active/ Inactive
INSAT-1A	April 10, 1982	Very High Resolution Radiometer(VHRR) Bands: Visible: 0.55-0.75µm IR : 10.5-12.5µm	Monitoring cyclones & monsoon CMV Winds OLR Rainfall Estimation	Inactive
INSAT-2B	August 8, 1983	do	do	Inactive
INSAT-1C	July 22, 1988	do	do	Inactive
INSAT-1D	June 12, 1990	do	do	Inactive

Satellite	Lunch date	Met Payload with Wavelength Bands	Major Applications	Active/ Inactive
INSAT-2A	July 10, 1992	Very High Resolution Radiometer(VHRR) Bands: 0.55-0.75µm 10.5-12.5µm	Monitoring cyclones & monsoon CMV Winds OLR Rainfall Estimation Mesoscale features Flood/Intense precipitation advisory Snow detection	Inactive
INSAT-2B	July 23, 1993	1. VHRR : payload Band : 0.55 – 0.75μm Vis: 2km 10.5 – 12.5μm IR: 8 km	do	Inactive
INSAT-2E	April, 1999	1.VHRR: As above +WV Band: 5.7 – 7.1μm 8km 2. CCD Payload Bands : 0.63 -0.79μm Vis: 2 km 0.77-0.86μm NIR 1.55-1.70μm SW IR: 1km	do	Active

Satellite	Lunch date	Met Payload with Wavelength Bands	Major Applications	Active/ Inactive
Kalpana-1	12th Sept., 2002	Very High Resolution Radiometer(VHRR) Bands: 0.55-0.75µm vis: 2km 10.5-12.5µm IR: 8 km WV Band: 5.7-7.1µm – 8km	Monitoring cyclones & monsoon CMV Winds OLR Rainfall Estimation	Active
INSAT-3A	10 April 2003	1. VHRR : payload Band : 0.55 – 0.75μm 10.5 – 12.5μm 5.7 – 7.1μm 2. CCD Payload Bands : 0.63 -0.68μm 0.77-0.86μm 1.55-1.70μm	As above & Mesoscale features Flood/intense precipitation advisory Snow detection	Active

# **FUTURE SATELLTE PROGRAM**

INSAT-3D	around 2006	6 channel Imager Band : 0.55 – 0.75μm (Vis- 1 km), 1.55 – 1.70μm (IR- 1km), 3.70-3.95μm(Mid Wave IR- 4 km), 6.5-7.10 μm (Thermal IR- 4 km), 10.3- 11.3μm (T IR- 4 km), 11.3- 12.5μm(WV- 8 km)	Monitoring cyclones & monsoon CMV Winds OLR Rainfall Estimation	
		Sounder Bands : 19 channels between 0.69-14.71 μm	Temperature and Humidity profiles in the atmosphere.	
Megha-Tropiques (A joint project by ISRO and CNES, France with the objective of studying the water cycle and	around 2006	SAPHIR 6 bands around 183 GHz (10 km Res.) SCARAB Radiation instrument in short & long wave (40 km Res.)	Water vapor profile upto 12 km Radiation budget	
energy exchanges in the tropics).		MADRAS 89 & 157 GHz radiometer 10, 18 & 37 GHz radiometer (10 km Res.)	Ice particles in cloud tops, cloud liquid water and precipitation; sea surface wind speed. 23 GHz : Integrated water vapor	

### CYCLONE WARNING DISSEMINATION SYSTEM (CWDS)

IMD has installed 250 specially designed receivers within the vulnerable coastal areas for direct transmission of warnings to the officials and people using broadcast capability of INSAT satellite. Recently this technology has been upgraded to digital transmission. Initially 100 Digital CWDS have been deployed in the coastal areas of Andhra Pradesh.

Area Cyclone Warning Center (ACWC) special warning bulletins transmitted every hour in local language to the affected areas.

### **METEOROLOGICAL DATA DISSEMINATION (MDD)**

Under this program analogue type cloud imagery data transmitted through INSAT S-band broadcast along with other conventional meteorological data and FAX analysed and forecast weather charts.

There are 33 MDD receiving stations in India and also operating in neighboring countries at Sri Lanka and Male under bi-lateral agreement. In general processed images are sent every three hous, and every hour during the cyclone period.

#### INTERNET PRODUCTS

INSAT satellite processed Global and Sector images (IR, Visible, WV) are provided through IMD Internet web (<u>www.imd.ernet.in</u>) every three hours, and every hour during the cyclone period in real-time.



### **OTHER SATELLITE PRODUCTS**

**PDUS for METEOSAT-5 data reception** 

A PDUS receiving station installed in early 2000 in IMD, New Delhi to receive high-resolution imagery data from METEOSAT-5 satellite located at 63<sup>0</sup> E.

### **RECEPTION OF NOAA SATELLITE DATA**

The AVHRR and TOVS data from NOAA series of polar orbiting satellites are received and processed by IMD at New Delhi and Chennai trough HRPT receiving station. The New Delhi HRPT receiving station upgraded to receive the NOAA (K, L, M, N series) satellite data.

#### **INDO-US co-operation**

INDO-US data exchange center has been established in 1999 for exchange of data with NOAA/NASA under the co-operation in Earth Atmospheric Sciences.

IMD providing INSAT cloud imagery data every three hours in real-time to NOAA/NASA and receiving GOES satellite series data.

These data products are being used by a number of scientists for specific studies.

### Satellite Products generated at IMD

- Cloud Motion Vectors (CMVs) Daily at 00 UTC processed operationally.
- Quantitative Precipitation Estimates (QPEs) based on modified Arkin method using cloud top temperature accumulated over 1x1 lat./long grid and correlated to rainfall.
- Outgoing Long-wave Radiation (OLR) is derived from thermal IR data using physical and statistical algorithms, based on radiative transfer principles.
- Sea Surface Temperatures (SST) is derived using the single channel (10.5-12.5) brightness temperature.
- Water vapor channel data (5.7-7.1) provides information on mid-tropospheric water vapor and flow pattern associated with incursion of water vapor during monsoon onset.
- Vertical Temperature Profiles (VTPRs)



<sup>(</sup>Background does not depict political boundary)

## **Synoptic Applications in IMD**

- Major application, the monitoring of Synoptic weather systems
- Watch and monitor growth of rapidly developing weather phenomena like cumulonimbus cells, thunderstorm complexes
- Identify and locate primary synoptic systems like troughs/ridges, jet streams and regions of intense convection, inter tropical convergence zones etc.
- Monitor onset and progress of monsoon
- Detect genesis and growth of Tropical Cyclones and monitor their intensification and movement till landfall.







# Future missions

- Several satellite missions have been planned to support the operational data needs and ongoing research efforts.
- The future INSAT missions will carry improved VHRR and vertical sounders for temperature/humidity profiles.
- The Megha -Tropiques Mission scheduled for 2006 with the objective of studying the water cycle and energy exchanges in the tropics.
- With an equatorial inclined orbit, the satellite will have high repeativity over tropical regions.
- The future appears bright for our space-based observing system. Advanced, multispectral (visible, IR, and passive microwave) imagers, sounders (infrared and microwave) and scatterometers are planned for launch in the near future.

- The satellite data downloads are expected to exceed several terabytes per day.
- Fortunately, communications and computing capacity are increasing at a rate that hopefully can accommodate this data explosion.
- It is important that the evolving space-based observational system keep one step ahead of the demands being placed by the user community and advances in numerical weather prediction.
- \* While it will become an enormous task and challenge to assimilate this wealth of data into meaningful parameters, the outlook is bright for unlocking the stillunresolved mysteries towards improving our understanding and prediction of atmospheric circulation systems such as tropical cyclones.



International TOVS Study Conference, 13<sup>th</sup>, TOVS-13, Sainte Adele, Quebec, Canada, 29 October-4 November 2003. Madison, WI, University of Wisconsin-Madison, Space Science and Engineering Center, Cooperative Institute for Meteorological Satellite Studies, 2003.