

Progress in modeling efforts related to radiance assimilation of water vapor, clouds, and precipitation

Ralf Bennartz¹, Tom Greenwald², Andrew Heidinger³,
Chris O'Dell¹, Martin Stengel^{1,6}, Peter Bauer⁵

1: Atmos. & Oceanic Sci., University of Wisconsin

2: CIMSS, University of Wisconsin

3: NOAA/NESDIS

4: NOAA/NCEP

5: ECMWF

6: SMHI


Outline

- Introduction
 - The SOI Model - Status
 - Water vapor from SEVIRI
 - Cloud/precipitation overlap parameterization
 - Future plans
-


The Successive Order of Interaction (SOI) radiative transfer model

- Fast forward IR and MW RT models (SOI) developed and tested.
 - Tangent linear and adjoint model developed and tested.
 - Model fully integrated in JCSDA CRTM
 - Application examples:
 - Bias monitoring and water vapor retrieval in IR
 - MW precipitation/cloud overlap parameterization for NWP
-

Infrared: Global SEVIRI IR bias monitoring against NCEP/GFS



Infrared: Global SEVIRI IR bias monitoring against NCEP/GFS




Optimal estimation of water vapor from SEVIRI

- Derives integrated water vapor, layer water vapor, and surface temperature
 - From SEVIRI window and WV channels
 - Integrated into EUMETSAT Climate-SAF
-


Optimal estimation of water vapor from SEVIRI

Water vapor column amount over water compared to passive microwave AMSR-E



Optimal estimation of water vapor from SEVIRI

Water vapor column amount over water compared to passive microwave AMSR-E



After 0.25 K bias correction in 8.7 micron channel

Optimal estimation of water vapor from SEVIRI

- Slight bias of window channel correction changes BIAS in WV retrieval from -2.5 kg/m^2 to 0.4 kg/m^2
 - Absolute calibration and spectral dependency (differences of window channels are giving most information about water vapor column)
 - High demands on spectral accuracy of surface emissivity. Land surfaces.....
 - We need to think about better ways to integrate LSE into retrieval
-


Forward modeling errors in microwave radiance assimilation of clouds/precipitation

- Uncertainties in gas absorption properties
 - Radiative transfer solver
 - Higher frequencies: Scattering of ice particles
 - 3D/beam filling errors
 - Handling of cloud overlap in forward model
-

Different cloud/precipitation overlap models

- **Conventional approach uses cloud cover to subdivide NWP pixel in cloudy/precipitation**
 - **New approach derives two/three optimal columns based on subscale distribution of precipitation columns with similar optical properties**
 - **Numerically efficient (2-3 radiative transfer calculations per NWP grid point)**
 - **Highly accurate against independent column/MR-overlap reference**
 - **Optimal approach reduces errors due to cloud overlap from maximum values of 5-10 K to values $< 1K$**
-


Different cloud/precipitation overlap models



**Currently
operational at
ECMWF**

O'Dell, Bauer, Bennartz,
JAS, 2006

Different cloud/precipitation overlap models



**Currently
operational at
ECMWF**

**New scheme with
much better error
characteristics**

O'Dell, Bauer, Bennartz,
JAS, 2006

Outlook

- Further development/refinement and integration of SOI into CRTM.
 - Address errors/uncertainties/biases associated with various cloud microphysics parameterizations in different numerical models
 - Expand work in infrared to infrared/application to AIRS.
-

International TOVS Study Conference, 15th, ITSC-15, Maratea, Italy, 4-10 October 2006
Madison, WI, University of Wisconsin-Madison, Space Science and Engineering Center,
Cooperative Institute for Meteorological Satellite Studies, 2006.