




*Using Cloudy  
AIRS Fields of  
View in  
Numerical  
Weather  
Prediction*



*John Le Marshall  
Director, JCSDA 2004-2007  
BoM 2007*



*J. Le Marshall*

*J. Jung*

*L.-P. Riishojgaard*

*M. Goldberg*

*C. Barnet*


*W. Wolf*

*J. Derber*

*R. Treadon*

*S. Lord*

.....



*John Le Marshall*  
*Director, JCSDA 2004-2007*  
*BoM 2007*



# Overview


---

- The Challenge
- CAWCR
- Recent Data Impact Studies
- Use of hyperspectral radiances in NWP – Cloudy Radiances
- Plans/Future Prospects
- Summary



# The Challenge Satellite Systems/Global Measurements

GRACE



Cloudsat




CALIPSO



Aqua



GIFTS




NPP



Landsat



GOES-R



NPOESS

NOAA/  
POES



SeaWiFS



Jason



ICESat



SORCE



Aura



WindSAT


TRMM



TOPEX  
SYSTEM



Meteor/  
SAGE



COSMIC/GPS



Terra



MSG





SSMIS




# 5-Order Magnitude Increase in Satellite Data Over 10 Years




## Daily Upper Air Observation Count



## Satellite Instruments by Platform




# Data Assimilation Impacts in the NCEP GDAS



Satellite and Conventional data provide nearly the same amount of improvement to the Northern Hemisphere.

**S. Hemisphere 1000 mb AC Z**  
**20S - 80S Waves 1-20**  
**15 Aug - 20 Sep '03**



**Anomaly correlation for days 0 to 7 for 500 hPa geopotential height in the zonal band 20°-80° for January/February. The red arrow indicate use of satellite data in the forecast model has doubled the length of a useful forecast.**



Australian Government  
Bureau of Meteorology




**CAWCR**


**The Centre for Australian  
Weather  
and Climate Research**





# CAWCR Partners








# Atmosphere-Land Observation & Assessment

- ❖ Atmospheric composition
  - gases
  - aerosol
- ❖ Cloud, radiation and precipitation processes
- ❖ Biogeochemical cycles (carbon & water)
- ❖ Micrometeorology
- ❖ Observing system technologies
- ❖ Remote sensing and data assimilation




Centre for Australian Weather and Climate Research: a partnership between CSIRO and the Bureau of Meteorology



Australian Government  
Bureau of Meteorology

# Scope of the Australian Community Climate and Earth System Simulator (ACCESS)



Centre for Australian Weather and Climate Research: a partnership between CSIRO and the Bureau of Meteorology





**SOME RECENT**

**ADVANCES / DATA IMPACT**



# *OBSERVING SYSTEM EXPERIMENTS*

## **OBSERVING SYSTEM EXPERIMENT WITH SATELLITE AND CONVENTIONAL DATA**

T. Zapotocny, J. Jung, J. Le Marshall, R Treadon, .....



The analysis and forecast model used for these observing system experiments is the NCEP Global Data Assimilation/Forecast System (GDAS/GFS).

The OSE consists of 45-day periods during January-February and August-September 2003. During these periods, a T254 - 64 layer version of NCEP's global spectral model was used.

The control run utilizes NCEP's operational data base and consists of all data types routinely assimilated in the GDAS. The two experimental runs have either all the conventional in-situ data denied (NoCon) or all the remotely sensed satellite data denied (NoSat). Differences between the control and experimental runs are accumulated over the 45-day periods and analyzed to demonstrate the forecast impact of these data types through 168 hours.

Note: geographic distribution of impact also calculated



**Table 1. Conventional data denied within the NCEP Global Data Assimilation System for this study. Mass observations (temperature and moisture) are shown in the left hand column while wind observations are shown in the right hand column.**

<b>Rawinsonde temperature and humidity</b>	<b>Rawinsonde u and v</b>
<b>AIREP and PIREP aircraft temperatures</b>	<b>AIREP and PIREP aircraft u and v</b>
<b>ASDAR aircraft temperatures</b>	<b>ASDAR aircraft u and v</b>
<b>Flight-level reconnaissance and dropsonde temperature, humidity and station pressure</b>	<b>Flight-level reconnaissance and dropsonde u and v</b>
<b>MDCARS aircraft temperatures</b>	<b>MDCARS aircraft u and v</b>
<b>Surface marine ship, buoy and c-man temperature, humidity and station pressure</b>	<b>Surface marine ship, buoy and c-man u and v</b>
<b>Surface land synoptic and Metar temperature, humidity and station pressure</b>	<b>Surface land synoptic and metar u and v</b>
<b>Ship temperature, humidity and station pressure</b>	<b>Wind Profiler u and v</b>
	<b>NEXRAD Vertical Azimuth Display u and v</b>
	<b>Pibal u and v</b>







**Table 2. Satellite data denied within the NCEP Global Data Assimilation System for this study.**

<b>HIRS sounder radiances</b>	<b>SBUV ozone radiances</b>
<b>MSU radiances</b>	<b>QuikSCAT surface winds</b>
<b>AMSU-A radiances</b>	<b>GOES atmospheric motion vectors</b>
<b>AMSU-B radiances</b>	<b>GMS atmospheric motion vectors</b>
<b>GOES sounder radiances</b>	<b>METEOSAT atmospheric motion vectors</b>
<b>SSM/I precipitation rate</b>	<b>SSM/I surface wind speed</b>
<b>TRMM precipitation rate</b>	













Fig. 6. Anomaly correlation for days 0 to 7 for 500 hPa geopotential height in the zonal band 20°-80° for each Hemisphere and season. The control simulation is shown in blue, while the NoSat and NoCon denial experiments are shown in magenta and green, respectively.

**a****N. Hemisphere 500 hPa AC Z 60N - 90N  
Waves 1-20 15 Jan - 15 Feb 2003****c****N. Hemisphere 500 hPa AC Z 60N - 90N  
Waves 1-20 15 Aug - 20 Sep 2003****b****S. Hemisphere 500 hPa AC Z 60S - 90S  
Waves 1-20 15 Jan - 15 Feb 2003****d****S. Hemisphere 500 hPa AC Z 60S - 90S  
Waves 1-20 15 Aug - 20 Sep 2003**

Anomaly correlation for days 0 to 7 for 500 hPa geopotential height in the polar cap region (60°-90°) of each Hemisphere and season. The control simulation is shown in blue, while the NoSat and NoCon denial experiments are shown in magenta and green, respectively.





Fig. 7 The impact of removing satellite and in-situ data on hurricane track forecasts in the GFS during the period 15 August to 20 September 2003. Panels (a and b) show the average track error (NM) out to 96 hours for the control experiment and the NoSat and NoCon denials for the Atlantic and Pacific Basins, respectively.



# *OBSERVING SYSTEM EXPERIMENTS*

## **OBSERVING SYSTEM EXPERIMENT WITH FOUR SATELLITE DATA TYPES AND RAWINSONDE DATA**


T. Zapotocny, J. Jung, J. Le Marshall, R Treadon, .....




A series of Observing System Experiments (OSEs) covering two seasons has been undertaken to quantify the contributions to the forecast quality from conventional rawinsonde data and from four types of remotely sensed satellite data.

The impact was measured by comparing the analysis and forecast results from an assimilation/forecast system using all data types in NCEP's operational data base with those from a system excluding a particular observing system.

For these OSEs, the forecast results are compared through 168 hours for periods covering more than a month during two seasons.





**Fig. 8** The day 5 anomaly correlations for waves 1-20 for the (a and d) mid-latitudes, (b and e) polar regions and (c and f) tropics. Experiments shown for each term include, from left to right, the control simulation and denials of AMSU, HIRS, GEO winds, Rawinsondes and QuikSCAT. The 15 January to 15 February 2003 results are shown in the left column and the 15 August to 20 September results are shown in the right column. Note the different vertical scale in (c and f).



**Fig. 9. The 15 January to 15 February 2003 day 0-7 500 hPa geopotential height die-off curves for the control and five denial experiments. The Northern Hemisphere results are shown in the left panels and the Southern Hemisphere results are shown in the right panels.**





**Fig. 10. Average track error (NM) by forecast hour for the control simulation and experiments where AMSU, HIRS, GEO winds and QuikSCAT were denied. The Atlantic Basin results are shown in (a), and the Eastern Pacific Basin results are shown in (b). A small sample size in the number of hurricanes precludes presenting the 96 hour results in the Eastern Pacific Ocean.**



# *OBSERVING SYSTEM EXPERIMENTS*

## **OBSERVING SYSTEM EXPERIMENT WITH NOAA POLAR ORBITING SATELLITES**

J. Jung, T. Zapotocny, J. Le Marshall, R Treadon, .....




An Observing System Experiments (OSEs) during two seasons has been used to quantify the contributions made to forecast quality from the use of the National Oceanic and Atmospheric Administration's (NOAA) polar orbiting satellites.


The impact is measured by comparing the analysis and forecast results from an assimilation/forecast system using observations from one NOAA polar orbiting satellite, NOAA-17 (1\_NOAA), with results from systems using observations from two, NOAA-16 and NOAA-17 (2\_NOAA), and three, NOAA-15, 16 and 17 (3\_NOAA), polar orbiting satellites.




(a) 0000 UTC 25 January 2003 NOAA 17




(b) 0000 UTC 25 January 2003 NOAA 16 and 17



(c) 0000 UTC 25 January 2003 NOAA 15, 16 and 17





**Fig. 12.** The day 5 anomaly correlations for waves 1-20 for the (a and d) mid-latitudes, (b and e) polar regions and (c and f) tropics. Experiments include data from 3\_NOAA, 2\_NOAA, and 1\_NOAA satellite(s). The 15 January to 15 February 2003 results are shown in the left column and the 15 August to 20 September 2003 results are shown in the right column. Note the different vertical scale in (c and f).





Fig. 15. Geographic distribution of Forecast Impact to 850 hPa relative humidity from the 2\_NOAA and 3\_NOAA experiments during August-September 2003. The 12, 24, 48 and 72-hr impacts are shown for each time period with the color contour interval 12.5%. Values within 12.5% of zero are white. Regions underground are shaded black.



*Advanced Sounders*

**Table 2.4-1 Characteristics of Advanced Infrared Sounders**

Name	AIRS	IASI	CrIS	IRFS	GIFTS
Orbit	705 km	833 km	824 km	1000 km	Geostationary
Instrument type	Grating	FTS	FTS	FTS	FTS
Agency and Producer	NASA JPL/LoMIRIS	EUMETSAT/ CNES Alcatel	IPO (DoD/NOAA/ NASA) ITT	Russian Aviation and Space Agency	NASA/NOAA/ Navy. Space Dynamics Lab.
Spectral range (cm <sup>-1</sup> )	649 –1135 1217–1613 2169 –2674	Contiguous 645-2760	650 -1095 1210 –1750 2155 –2550	625 -2000 2200 -5000	685-1130 1650-2250
Unapodized spectral resolving power	1000 – 1400	2000 – 4000	900 – 1800	1200 - 4000	2000-6000
Field of view (km)	13 x 7	12	14	20	4
Sampling density per 50 km square	9	4	9	1	50
Power (W)	225	200	86	120	254
Mass (kg)	140	230	81	70	59
Platform	AQUA (EOS PM1)	METOP-1,-2,-3	NPP and NPOESS C1	METEOR 3MN2	Geostationary
Launch date	Feb 2002	2006	2010 for NPP 2013 NPOESS C1	2010+	2010+?






# *AIRS Data Assimilation*

*J. Le Marshall, J. Jung, J. Derber, R. Treadon,*

*S.J. Lord, M. Goldberg, C. Barnet, W. Wolf and H-S Liu, J. Joiner,*

*and J Woollen.....*



**1 January 2004 – 31 January 2004**

**Used operational GFS system as Control**

**Used Operational GFS system Plus AIRS  
as Experimental System**

# Background




- Atmospheric Infrared Sounder (AIRS) was launched on the AQUA satellite on May 4, 2002 - Polar orbit 705 km, 13:30 ECT
- AIRS – high spectral resolution infrared sounder, demonstrated significantly improved accuracy of temperature and moisture soundings.
- NOAA/NESDIS is processing and distributing AIRS data and products in near real-time to operational NWP centers.





# AIRS IR Instrument



- AIRS is a cooled grating array spectrometer
- Spectral coverage 3.7 to 15.4 microns in 17 arrays with 2378 spectral channels (3.74-4.61  $\mu\text{m}$ , 6.2-8.22  $\mu\text{m}$ , 8.8-15.4  $\mu\text{m}$ )
- Spectral resolution  $\lambda/\Delta\lambda=1200$ , 14 km FOV from 705km orbit
- Launch – May 2002
- Primary products: temperature profile (< 1 K accuracy), moisture profile (< 15%), ozone (< 15 % (layers) and 3 % total)
- Research products: CO<sub>2</sub>, CO, CH<sub>4</sub>
- The integrated sounder system includes the AIRS VIS/NIR channels and microwave sounders

*Table 1: Satellite data used operationally within the NCEP Global Forecast System*


<ul style="list-style-type: none"><li><b>HIRS sounder radiances</b></li><li><b>AMSU-A sounder radiances</b></li><li><b>AMSU-B sounder radiances</b></li><li><b>GOES sounder radiances</b></li><li><b>GOES 9,10,12, Meteosat atmospheric motion vectors</b></li><li><b>GOES precipitation rate</b></li><li><b>SSM/I ocean surface wind speeds</b></li><li><b>SSM/I precipitation rates</b></li></ul>	<ul style="list-style-type: none"><li><b>TRMM precipitation rates</b></li><li><b>ERS-2 ocean surface wind vectors</b></li><li><b>Quikscat ocean surface wind vectors</b></li><li><b>AVHRR SST</b></li><li><b>AVHRR vegetation fraction</b></li><li><b>AVHRR surface type</b></li><li><b>Multi-satellite snow cover</b></li><li><b>Multi-satellite sea ice</b></li><li><b>SBUV/2 ozone profile and total ozone</b></li></ul>
---	---

# *Global Forecast System Background*


- Operational SSI (3DVAR) version used
- Operational GFS T254L64 with reductions in resolution at 84 (T170L42) and 180 (T126L28) hours. 2.5hr cut off



AQUA AIRS 20040131 06Z  
Observed-Calculated Brightness Temperature with Bias Correction



Channel 051 Freq 661.8 cm<sup>-1</sup> Nobs 7070 Avg. 0.038 Std. 0.73




**AIRS data coverage at 06 UTC on 31 January 2004. (Obs-Calc. Brightness Temperatures at 661.8 cm<sup>-1</sup> are shown)**

*Table 2: AIRS Data Usage per Six Hourly Analysis Cycle*

Data Category	Number of AIRS Channels
Total Data Input to Analysis	~200x10 <sup>6</sup> radiances (channels)
Data Selected for Possible Use	~2.1x10 <sup>6</sup> radiances (channels)
Data Used in 3D VAR Analysis(Clear Radiances)	~0.85x10 <sup>6</sup> radiances (channels)


**S. Hemisphere 1000 mb AC Z**  
**20S - 80S Waves 1-20**  
**1 Jan - 27 Jan '04**



**Figure1(a). 1000hPa Anomaly Correlations for the GFS with (Ops.+AIRS) and without (Ops.) AIRS data, Southern hemisphere, January 2004**



**N. Hemisphere 1000 mb AC Z  
20N - 80N Waves 1-20  
1 Jan - 27 Jan '04**




**Figure3(a). 1000hPa Anomaly Correlations for the GFS with (Ops.+AIRS) and without (Ops.) AIRS data, Northern hemisphere, January 2004**



## AIRS Data Assimilation

J. Le Marshall, J. Jung, J. Derber, R. Treadon, S.J. Lord,  
M. Goldberg, W. Wolf and H-S Liu, J. Joiner and J Woollen




**January 2004**

**Used operational GFS system as Control**

**Used Operational GFS system Plus AIRS  
as Experimental System**

**Clear Positive Impact Both Hemispheres. Implemented -2005**





# *AIRS Data Assimilation*

## *Impact of Data density...*

**10 August – 20 September 2004**

**N. Hemisphere 500 mb AC Z**  
**20N - 80N Waves 1-20**  
**10 Aug - 20 Sep '04**






# *AIRS Data Assimilation*

## *Impact of Spectral coverage...*

**10 August – 20 September 2004**

**Day 5 Average Anomaly Correlation**  
**Waves 1- 20**  
**2 Jan - 15 Feb 2004**





# *AIRS Data Assimilation*


## *MOISTURE*

*Forecast Impact* evaluates which forecast (with or without AIRS) is closer to the analysis valid at the same time.


$$\text{Impact} = 100 * [\text{Err}(\text{Cntl}) - \text{Err}(\text{AIRS})] / \text{Err}(\text{Cntl})$$

Where the first term on the right is the error in the Cntl forecast. The second term is the error in the AIRS forecast. Dividing by the error in the control forecast and multiplying by 100 normalizes the results and provides a percent improvement/degradation. A positive Forecast Impact means the forecast is better with AIRS included.

AIRSC 024-HR 925 hPa RH Fcst Imp (%) (15 Jan-15 Feb 2004)







# *AIRS Data Assimilation*


## *Using Cloudy Fields of View*

**1 January – 24 February 2007**



# *AIRS Data Assimilation*

## *Using Cloudy Fields of View*



**Initial Experiments: 1 January – 24 February 2007**

**Intention:**

**Assimilate radiances from cloudy fofs preferably with single level cloud.**

**Initially use radiances where cloud coverage and uniformity of fofs allow accurate estimation of radiances from clear part of fofs**


**Initially measure impact from use of clear air radiances**

**(Later use  $\alpha$  and  $p_c$  in 3D Var.)**




# *AIRS Data Assimilation*

## *Using Cloudy Fields of View*



**Initial approach to use 9 AIRS fofs and AMSU-A data to provide cloud level information and provide error characterized radiances from clear part of fof.**

**Subsequently MODIS information to be used as well to improve cloud characterization (ensure single level cloud) and provide error characterized radiances from clear part of fof, cloud height and cloud amount.**



# *AIRS Data Assimilation*

## *Using Cloudy Fields of View*

**Initial Experiments: 1 January – 24 February 2007**

**Assume :**

$$R_j = (1 - \alpha_j) R_{\text{clr}} + \alpha_j R_{\text{cld}}$$


**Only variability in AIRS fov is cloud amount  $\alpha_j$**

**9 AIRS fovs on each AMSU-A footprint used to estimate  $R_{\text{clr}}$**




# *AIRS Data Assimilation*

## *Using Cloudy Fields of View*



**Susskind, J., C.D. Barnet and J.M. Blaisdell 2003. Retrieval of atmospheric and surface parameters from AIRS/AMSU/HSB data in the presence of clouds. IEEE Trans. Geosci. Remote Sens., 41, 390-409.**



# *AIRS Data Assimilation*

## *Using Cloudy Fields of View*


**Initial Experiments: 1 January – 24 February 2007**

**Control – Current Ops. ( OP. data coverage - Uses 152 AIRS channels from all fofs with operational thinning)**

**Experiment- Op. data coverage, minus Op. AIRS plus AIRS radiances from channels free from cloud effects and radiances from the clear air part of selected cloudy fofs (with operational thinning).**




### N. Hemisphere 500 hPa AC Z 20N - 80N Waves 1-20 1 Jan - 24 Feb '04



— Control — AIRS (cloudy fofs)



### S. Hemisphere 500 hPa AC Z 20S - 80S Waves 1-20 1 Jan - 24 Feb '07








# *AIRS Data Assimilation*

## *Using Cloudy Fields of View*



**Initial Experiments: 1 January – 24 February 2007**

### **Results:**

**Assimilation of radiances from cloudy fovs resulted in improved anomaly correlations for the experimental system during the period studied.**

**Southern Hemisphere results significant at near the 95% level, accounting for serial correlation of forecast differences (Seaman, 1992)**

**Further R2O activity restricted by loss of RT data set.**



# Surface Emissivity ( $\epsilon$ ) Estimation

Emissivity ( $\epsilon$ ) required for

- Accurate surface temperature
- Accurate Boundary layer temperature
- Accurate Boundary layer moisture




# Surface Emissivity ( $\epsilon$ ) Estimation Methods

- Geographic Look Up Tables (LUTs) - CRTM
- Regression based on theoretical estimates
  - Lihang Zhou
- Minimum Variance, provides  $T_{\text{surf}}$  and  $\epsilon^*$
- Eigenvector technique
  - Dan Zhou and Bill Smith
- Variational Minimisation – goal

# Regression IR HYPERSENSPECTRAL EMISSIVITY - ICE and SNOW

Sample Max/Min Mean computed from synthetic radiance sample



From Lihang Zhou

# Surface Emissivity ( $\epsilon$ ) Estimation Methods

## JCSDA IR Sea Surface Emissivity Model (IRSSE)

Initial NCEP IRSSE Model based on Masuda et al. (1998)

Updated to calculate Sea Surface Emissivities via Wu and Smith (1997)

Van Delst and Wu (2000)

Includes high spectral resolution (for instruments such as AIRS)

Includes sea surface reflection for larger angles

JCSDA Infrared Sea Surface Emissivity Model – Paul Van Delst  
Proceedings of the 13th International TOVS Study Conference  
Ste. Adele, Canada, 29 October - 4 November 2003



## AIRS SST and $\epsilon$ Determination

Use AIRS bias corrected radiances from GSI

AIRS channels used are :

119 – 129 (11)

154 – 167 (14)

263 – 281 (19)

Method is the minimum (emissivity) variance technique

Channels used in Pairs : 119, 120; 120, 121; 121, 122; .. etc

For a downward looking infrared sensor:

$$I_{\nu} = \int_0^Z B_{\nu}[T(z)] \frac{\partial \tau_{\nu}(z, Z)}{\partial z} dz + \varepsilon_{\nu} \bullet B_{\nu}(T_S) \bullet \tau_{\nu}(0, Z) +$$

$$(1 - \varepsilon_{\nu}) \bullet \tau_{\nu}(0, Z) \int_{\infty}^0 B_{\nu}[T(z)] \frac{\partial \tau_{\nu}(z, Z)}{\partial z} dz$$

where  $I_{\nu}$ ,  $\varepsilon_{\nu}$ ,  $B_{\nu}$ ,  $T_S$ ,  $\tau_{\nu}(z_1, z_2)$ ,  $Z$  and  $T(z)$  are observed spectral radiance, spectral emissivity, spectral Planck function, the surface temperature, spectral transmittance at wavenumber  $\nu$  from altitude  $z_1$  to  $z_2$ , sensor altitude  $z$ , and air temperature at altitude  $z$  respectively.

The solution can be written as :

$$\hat{\varepsilon}_v = \frac{[R_v^{OBS} - N_v^\uparrow] - \tau_v \overline{N}_v^\downarrow}{\tau_v B_v(\hat{T}_S) - \tau_v \overline{N}_v^\downarrow}$$


Where  $R^{OBS}$  is the observed upwelling radiance,  $N^\uparrow$  represents the upwelling emission from the atmosphere only and  $N^\downarrow$  represents the downwelling flux at the surface. The  $\hat{\phantom{x}}$  symbol denotes the “effective” quantities as defined in Knuteson et al. (2003).

The SST is the  $T_S$  that minimises :

$$\sum (\varepsilon_i - \varepsilon_{i+1})^2$$




January 2007






# Ocean Surface Emissivity Comparisons



**Average surface emissivity over ocean as derived from AIRS using the minimum variance (AIRS Derived) and values from the ocean emissivity model within the CRTM (CRTM Calculated).**




# Ocean Emissivity Scan Angle Comparisons



**Average surface emissivity over ocean by scan angle as derived from AIRS using the minimum variance (AIRS Calculated) and values from the ocean emissivity model within the CRTM (CRTM Calculated).**




# Surface Emissivity Comparisons for Snow



Comparison of surface emissivity for snow from the CRTM lookup table (CRTM\_emissivity) with the emissivity derived from AIRS using minimum variance (AIRS\_derived\_emissivity)




# Surface Emissivity Comparisons for Ice



**Comparison of surface emissivity for ice from the CRTM lookup table (CRTM\_emissivity) with the emissivity derived from AIRS using minimum variance (AIRS\_derived\_emissivity).**



# Surface Emissivity Comparisons for Pine Forest



Comparison of surface emissivity for Pine Forest from the CRTM lookup table (CRTM\_emissivity) with the emissivity derived from AIRS using minimum variance (AIRS\_derived\_emissivity)



# The Future

---

- Expanded use of the hyperspectral - IASI
- Expanded use of cloudy radiances
- Improved surface emissivity characterization
- Expanded use of spectral content
- Enhanced use of moisture information
- ACCESS – UKUM

Use of Continuous Data in 4D-VAR (Regional 37.5km )


eg. TC Nicholas Western Australian region February  
2008

- .....

# The Future



**N. Hemisphere 500 hPa AC Z  
20N - 80N Waves 1-20  
1 Dec 2007 - 12 Jan 2008**






# The Future



**S. Hemisphere 500 hPa AC Z**  
**20S - 80S Waves 1-20**  
**1 Dec 2007 - 12 Jan 2008**



— Control — IASI\_EUMETSAT



# The Future

---

- Expanded use of the hyperspectral - IASI
- Expanded use of cloudy radiances
- Improved surface emissivity characterization
- Expanded use of spectral content
- Enhanced use of moisture information
- ACCESS – UKUM


Use of Continuous Data in 4D-VAR (Regional 37.5km )

eg. TC Nicholas Western Australian region February 2008

- .....

*Satellite data used within the ACCESS (UKUM) Forecast System Includes*

<p><b>HIRS sounder radiances</b> <b>AMSU-A sounder radiances</b> <b>AMSU-B sounder radiances</b> <b>GOES 10,12, Meteosat, MTSat-1R</b> <b>atmospheric motion vectors</b> <b>SSM/I ocean surface wind speeds</b></p>	<p><b>ERS-2 ocean surface wind vectors</b> <b>Quikscat ocean surface wind vectors</b> <b>AIRS sounder radiances</b> <b>Local hourly AMVs</b> <b>.....</b></p>
---	---




WIND(MPS) PW=100 399 ERR=0 0


NICHOLAS 04-08UTC(VIS/IR AN

1 0001 GMS-5 2 15 FEB 08046 060000 05838 01576 04.00

Thursday 14 February 2008 12UTC MELBN Forecast +15 VT: Friday 15 February 2008 03UTC Surface: mean sea level pressure



Thursday 14 February 2008 12UTC MELBN Forecast +24 VT: Friday 15 February 2008 12UTC Surface: mean sea level pressure





# *Summary*

Key components of the operational data base have been assessed in terms global forecast impact.

Quantitative estimates (ACs, FIs and hurricane forecast track errors) have been used to quantify the impact of conventional data, satellite data, and that of particular instruments and rawinsonde data in a number of OSEs.

In these studies the significant impacts of AMSU and rawinsondes were noted.



## *Summary / Cont'd*

AIRS (hyperspectral radiance) data have been shown to make a very significant contribution globally to operational NWP.

The significant potential for larger benefits to operational meteorology from use of these hyperspectral radiances has also demonstrated:

Data impact studies showing the importance of using improved spatial and spectral resolution data and showing the benefit of using cloud effected radiances have been presented.

Assimilation studies with UKUM based ACCESS model underway and showing good forecast skill.




An aerial photograph showing a dense, intricate network of rivers and streams flowing through a mountainous landscape. The terrain is rugged, with steep slopes and numerous peaks. The rivers are depicted in various shades of blue and brown, contrasting with the green and brown vegetation on the surrounding land. The overall scene conveys a sense of a complex, interconnected water system.

**The business of looking down  
is looking up**



**Australian Government**

**Bureau of Meteorology**



**CSIRO**



International TOVS Study Conference, 16<sup>th</sup>, ITSC-16, Angra dos Reis, Brazil, 7-13 May 2008.  
Madison, WI, University of Wisconsin-Madison, Space Science and Engineering Center,  
Cooperative Institute for Meteorological Satellite Studies, 2008.