



The Assimilation of Clear-Sky Infrared Radiances in the HIRLAM model at SMHI

Martin Stengel

SMHI, Swedish Meteorological and Hydrological Institute

The Assimilation of Clear-Sky Infrared Radiances in the HIRLAM Model

Martin Stengel (martin.stengel@smhi.se)

SMHI, Swedish Meteorolog



The potential use of spatially and temporal high resolved SEVIRI observations in the time dimension in HIRLAM 4D-Var is appropriate to make full use of SEVIRI environment using the two water vapour channels (WV6.2,WV7.3), located at 4.2 μ m. The impact on the analysis are illustrated for one case study. Secondly, we present upper air parameters during the summer period. In the winter period, the impact on

Data Preparation

The SEVIRI observations undergo various data preparation steps which are listed below. Only 500 to 1200 pixels (two observations each) approx. 8000 pixels are then used in each observation window. (6 obs./windows per 40-yr analysis)

- Processing of BTs and PGs using the SAF NWC software (for SEVIRI segments 7 and 8)
- Selecting 1 pixel out of a 16x16 pixel box
- Rejecting out-of-domain pixels (and scan angles >= 70°)
- Rejecting cloudy pixels (POD1) / OM, cloud mask
- Applying flat bias correction to WV6.2 obs. (2.4K)
- First guess check
- Spatial thinning (thinning bascule = 92km)



As demonstrated the middle air can be seen here.



Fig. 3: Observations with assimilated

Current use of satellite data in HIRLAM/SMHI:

- AMSU-A
 - NOAA 15, NOAA 16
 - Channels 5 to 10
 - 4D-Var
- (Per Dahlgren)

The Assimilation of Clear-Sky Infrared Radiances in the HIRLAM Model


Martin Stengel (martin.stengel@smhi.se)

SMHI, Swedish Meteorological and Hydrological Institute



Keywords:

- HIRLAM/SMHI (limited area model)
- 4D-Var
- SEVIRI water vapour channels
(data from 6 time slots used)
- not operational yet



SEVIRI IR radiances have been introduced successfully as a new observation type in summer month and one winter month. With the chosen experimental specification, mid- and upper-tropospheric humidity fields. A positive impact can be seen in the with forecast length. Small positive effects are also found for the temperature, geopotential height and wind fields. The impact on the temperature, geopotential height and wind fields is found to be very small and rather neutral during the winter period. "Upgrading", will be addressed in the near future.

This poster is presenting research results with contributions from: Martin Stengel, SMHI and others. This work is supported by EUMETSAT within the EUMETSat Reanalysis program.

The Assimilation of Clear-Sky Infrared Radiances in the HIRLAM Model

Martin Stengel (martin.stengel@smhi.se)

SMHI, Swedish Meteorological and Hydrological Institute



Abstract

The potential use of spatially and temporal high resolved SEVIRI observations in the HIRLAM model 4D-Var analysis is currently investigated at SMHI. Especially, the extended utilization of the time dimension in HIRLAM is appropriate to make full use of SEVIRI's fast imaging cycle. Currently, we carry out observation impact studies in a pre-operational environment using the two water vapour channels (WV6.2,WV7.3), located at 6.2μm and 7.3μm, to find an optimal assimilation setup. Firstly, the data preparation steps as well as the impact on the analysis are illustrated for one case study. Secondly, we present on this poster two impact studies, which show a positive impact of SEVIRI observations on the upper air parameters during the summer period. In the winter period, the impact on moisture is less pronounced and rather neutral for the other variables.

Data Preparation

- The SEVIRI observations undergo various data preparation steps which are listed below. Usually 500 to 1200 pixels (two observations each) at approx. 90km resolution are then kept in each observation window. (6 obs.-windows per 4D-Var analysis)
- Processing of BTs and PGEs using the SAF NWC software (for SEVIRI-segments 7 and 8)
- Selecting 1 pixel out of a 10x10 pixel box
- Rejecting out-of-domain pixels (and scan angles .gt. 70°)
- Rejecting cloudy pixels (PGE01/CMa, cloud mask)
- Applying flat BIAS correction to WV6.2 obs. (2.6K)
- First guess check
- Spatial thinning (thinning box size = 90km)




Fig.1: Example of thinned SEVIRI observations (red dots) after applying all data preparation steps.





Fig.2: Spatial distribution of used SEVIRI pixels for a one month period with 4D-Var.

Impact on HIRLAM 4D-Var Analysis - Case Study



As demonstrated in this case study, the impact of using the middle and upper troposphere (figure 3). This large difference can be seen here, is due to a systematic deviation in the background fields.

Observation Impact Study – Summer Period

Experiment description

- Time period : 08/24/2005 - 07/23/2005
- Control run (CTRL4D) : 4D-Var analysis with conventional observations only
- Experiment (SEV4D) : 4D-Var analysis with conventional and SEVIRI observations; SEVIRI observations are taken from six time slots (the slot closest to the respective observation window centre)
- Cycle : 6 hour assimilation cycle / forecast up to 48 hours
- Verification : EUMETSat radiosondes and synops




Fig.6: Time series of all lower prognostic height forecast errors. Shown are daily means of total and bias.




Fig.7: Time series of all lower prognostic height forecast errors. Shown are daily means of total and bias.

Observation

- Experiment description
- Time period : 12/01
- Control run (CTRL4D) : 4D-Var with SEVIRI observations only
- Experiment (SEV4D) : 4D-Var with SEVIRI observations only
- Cycle : 6 hours
- Verification : EUMETSat radiosondes and synops



Fig.8: Time series of all lower prognostic height forecast errors. Shown are daily means of total and bias.




Fig.9: Time series of all lower prognostic height forecast errors. Shown are daily means of total and bias.

Summary

SEVIRI IR radiances have been introduced successfully as a new observation type in the HIRLAM assimilation system. After a summer month and one winter month, with the chosen experimental specifications, the assimilation of SEVIRI's water vapour mid- and upper-tropospheric fields. A positive impact can be seen in the upper troposphere for all experiments with forecast length. Small positive effects are also found for the temperature, geopotential height and wind fields after 48 h forecast. It is found to be very small and rather neutral during the winter period. Changes in the data preparation step "Averaging", will be addressed in the near future.

Acknowledgements

This poster is presenting research results with contributions from: Martin Stengel, Per Delgren, Magnus Lindberg, Per Under, SMHI and others. This work is supported by EUMETSAT within the EUMETSAT Research program.

Data Preparation

The SEVIRI observations undergo various data preparation steps which are listed below. Usually 500 to 1200 pixels (two observations each) at approx. 90km resolution are then kept in each observation window. (6 obs.-windows per 4D-Var analysis)

- Processing of BTs and PGEs using the SAF NWC software (for SEVIRI-segments 7 and 8)
- Selecting 1 pixel out of a 10x10 pixel box
- Rejecting out-of-domain pixels (and scan angles .gt. 70°)
- Rejecting cloudy pixels (PGE01/CMa, cloud mask)
- Applying flat BIAS correction to WV6.2 obs. (2.6K)
- First guess check
- Spatial thinning (thinning box size = 90km)




Fig.1: Example of thinned SEVIRI observations (red dots) after applying all data preparation steps.




Fig.2: Spatial distribution of used SEVIRI pixels for a one month period with 4D-Var.

The Assimilation of Clear-Sky Infrared Radiances in the HIRLAM Model

Martin Stengel (martin.stengel@smhi.se)

SMHI, Swedish Meteorological and Hydrological Institute




Abstract

The potential use of spatially and temporal high resolved SEVIRI observations in the HIRLAM model 4D-Var analysis is currently investigated at SMHI. Especially, the extended utilization of the time dimension in HIRLAM 4D-Var is appropriate to make full use of SEVIRI's fast imaging cycle. Currently, we carry out observation impact studies in a pre-operational environment using the two water vapour channels (WV6.2,WV7.3), located at 4.6μm and 7.3μm, to find an optimal assimilation setup. Firstly, the data preparation steps as well as the impact on the analysis are illustrated for one case study. Secondly, we present on this poster two impact studies, which show a positive impact of SEVIRI observations on the upper air parameters during the summer period. In the winter period, the impact on moisture is less pronounced and rather neutral for the other variables.

Data Preparation

The SEVIRI observations undergo various data preparation steps which are listed below. Only 500 to 1200 pixels (two observations each) approx. 8000 pixels are then used in each observation window. (6 obs./windows per 40-Year analysis)

- Processing of BTs and PGs using the SAF NWC software (for SEVIRI segments 7 and 8)
- Selecting 1 pixel out of a 16x16 pixel box
- Rejecting out-of-domain pixels (and scan angle > 70°)
- Rejecting cloudy pixels (PO201/OM4, cloud mask)
- Applying Net Bias correction to WV6.2 obs. (2.4K)
- First guess check
- Spatial thinning (thinning baseline = 95km)



Impact on HIRLAM 4D-Var Analysis - Case Study

As demonstrated in this case study, the impact of assimilated SEVIRI radiances is mainly in the moisture fields of the middle and upper troposphere (Figure 3). The large-scale drying in the upper-tropospheric layers, which can be seen here, is due to a systematic deviation in the background minus observation departures (Figure 4).

Impact on HIRLAM 4D-Var Analysis - Case Study

As demonstrated in this case study, the impact of assimilated SEVIRI radiances is mainly in the moisture fields of the middle and upper troposphere (Figure 3). The large-scale drying in the upper-tropospheric layers, which can be seen here, is due to a systematic deviation in the background minus observation departures (Figure 4).

Observation Impact Study I

Experiment description:

- Time period : 05/24/2005 - 07/23/2005
- Control run (CTR4D) : 4D-Var analysis with conventional observations only
- Experiment (SEV4D) : 4D-Var analysis with co-assimilated SEVIRI observations closest to the respective observation times
- Cycle : 6 hour assimilation cycle
- Verification : ENGLAN radiances as background







Fig.3: Increments in Relative Humidity at 3 atmospheric levels (700hPa:left, 500hPa:middle, 250hPa right), calculated with respect to a control analysis with conventional observations only.


Channel: WV6.2



Channel: WV7.3



Channel: WV6.2



Channel: WV7.3




Fig.4: Frequency distribution of background minus observation departures.

Fig.5: Frequency distribution of analysis minus observation departures.

SEVIRI IR radiances have been introduced successfully during the summer month and one winter month, with the middle and upper-tropospheric humidity fields. A gain in forecast length, small positive effects are also visible. It is found to be very small and rather "superficial", will be addressed in the near future.

This poster is presenting research results with contributions from SMHI and others. This work is supported by EU-EU-





Fig.11: RMSE and Bias (against radiosonde data) for upper air parameters for the control run (CTRL4D, red lines) and SEVIRI experiment (SEV4D, dashed green lines). First row: at analysis time; second row: after 48 hours forecast.



SEVIRI IR radiances have been introduced successfully as a new observation type in the NMC/NM model assimilation system. Assimilation experiments shown in this presentation cover one summer month and one winter month. With the chosen experimental specification, the assimilation of SEVIRI's water vapour channels primarily affects the analysed and forecasted mid- and upper-tropospheric humidity fields. A positive impact can be seen in the upper troposphere for all experiments when comparing to radiosondes. This impact is decreasing with forecast length. Small positive effects are also found for the temperature, geopotential height and wind fields after 48 hours in the summer period. The impact on those upper air variables is found to be very small and rather neutral during the winter period. Changes in the data preparation step, as for example better bias correction and possibly "superclouding", will be addressed in the near future.

Acknowledgments

This poster is presenting research results with contributions from: Martin Stengel, Per Delgren, Magnus Lindberg, Per Uden, Niels Gustafsson, Anke Thors, Tomas Landellus (all SMHI) and others. This work is supported by EUMETSAT within the EUMETSat Reference program.





Fig.8: RMSE and Bias (against radiosonde data) for upper air parameters for the control run (CTRL4D, red lines) and SEVIRI experiment (SEV4D, dashed green lines). First row: at analysis time; second row: after 48 hours forecast.



Summary
SEVIRI IR radiances have been introduced successfully as a new observation type in the NMC assimilation system. Assimilation experiments shown in this presentation cover one summer month and one winter month. With the chosen experimental specification, the assimilation of SEVIRI's water vapour channels primarily affects the analysed and forecasted mid- and upper-tropospheric fields. A positive impact can be seen in the upper troposphere for all experiments when comparing to radiosondes. This impact is decreasing with forecast length. Small positive effects are also found for the temperature, geopotential height and wind fields after 48 hours in the summer period. The impact on those upper air variables is found to be very small and rather neutral during the winter period. Changes in the data preparation step, as for example better bias correction and possibly "superclouding", will be addressed in the near future.

Acknowledgments
This poster is presenting research results with contributions from: Martin Stengel, Per Delgren, Magnus Lindberg, Per Uden, Niels Gustafsson, Anke Thors, Tomas Landellus (all SMHI) and others. This work is supported by EUMETSAT within the EUMETSAT framework program.

The Assimilation of Clear-Sky Infrared Radiances in the HIRLAM Model

Martin Stengel (martin.stengel@smhi.se)

SMHI, Swedish Meteorological and Hydrological Institute




Abstract

The potential use of spatially and temporal high resolved SEVIRI observations in the HIRLAM model 4D-Var analysis is currently investigated at SMHI. Especially, the extended utilization of the time dimension in HIRLAM is appropriate to make full use of SEVIRI's fast imaging cycle. Currently, we carry out observatory impact studies in a pre-operational environment using the two water vapour channels (WV6.2,WV7.3), located at 4.2 μ m and 7.3 μ m, to find an optimal assimilation setup. Firstly, the data preparation steps as well as the impact on the analysis are illustrated for one case study. Secondly, we present on this poster two impact studies, which show a positive impact of SEVIRI observations on the upper air parameters during the summer period. In the winter period, the impact on moisture is less pronounced and rather neutral for the other variables.

Data Preparation


The SEVIRI observations undergo various data preparation steps which are listed below. Usually 500 to 1200 pixels (two observations each) at approx. 80km resolution are then used. In this observation window, (6 obs./windows per 40-hr analysis)

- Processing of BTs and PGs using the SAF NWC software (for SEVIRI segments 7 and 6)
- Selecting 1 pixel out of a 16x16 pixel box
- Rejecting out-of-domain pixels (and scan angles > 70°)
- Rejecting cloudy pixels (POE01/Dm, cloud mask)
- Applying Net Bias correction to WV6.2 obs. (2.4K)
- First guess check
- Spatial thinning (thinning bascule = 95km)



Impact on HIRLAM 4D-Var Analysis - Case Study


As demonstrated in this case study, the impact of assimilated SEVIRI radiances is mainly in the moisture fields of the middle and upper troposphere (Figure 3). This large-scale drying in the upper-tropospheric layers, which can be seen here, is due to a systematical difference in the background minus observation departures (Figure 4).



Observation Impact Study I - Summer Period

Experiment description


- Time period : 05/24/2005 - 07/21/2005
- Control run (CTRL4D) : 4D-Var analysis with conventional observations only
- Experiment (SEV4D) : 4D-Var analysis with conventional and SEVIRI observations; SEVIRI observations are taken from six time slots (the slot closest to the respective observation window centre)
- Cycle : 6 hour assimilation cycle / forecast up to 48 hours
- Verification : ENGLAM radiosondes and synops



Observation Impact Study II - Winter Period

Experiment description

- Time period : 12/01/2005 - 12/31/2005
- Control run (CTRL4D) : 4D-Var analysis with conventional observations only
- Experiment (SEV4D) : 4D-Var analysis with conventional and SEVIRI observations; SEVIRI observations are taken from six time slots (the slot closest to the respective observation window centre)
- Cycle : 6 hour assimilation cycle / forecast up to 48 hours
- Verification : ENGLAM radiosondes and synops



Presentation number: 6.12

Summary

SEVIRI IR radiances have been introduced successfully as a new observation type in the HIRLAM assimilation system. Assimilation experiments shown in this presentation cover one summer month and one winter month. With the chosen experimental specifications, the assimilation of SEVIRI's water vapour channels primarily affects the analysed and forecasted mid- and upper-tropospheric fields. A positive impact can be seen in the upper troposphere for all experiments when comparing to radiosondes. This impact is decreasing with forecast length. Small positive effects are also found for the temperature, geopotential height and wind fields after 48 hours in the summer period. The impact on those upper air variables is found to be very small and rather neutral during the winter period. Changes in the data preparation steps, as for example better bias correction and possibly "superclouding", will be addressed in the near future.

Acknowledgments

This poster is presenting research results with contributions from: Martin Stengel, Per Dahlberg, Magnus Lindberg, Per Under, Håk Gustafsson, Anke Thors, Tomas Landellus (all SMHI) and others. This work is supported by EUMETSAT within the EUMETSat Fellowship program.

Thank you

International TOVS Study Conference, 16th, ITSC-16, Angra dos Reis, Brazil, 7-13 May 2008.
Madison, WI, University of Wisconsin-Madison, Space Science and Engineering Center,
Cooperative Institute for Meteorological Satellite Studies, 2008.